



# LYME DISEASE

1. **Agent:** *Borrelia burgdorferi*, a spirochete first identified in 1982.

2. **Identification:**

a. **Symptoms:** Lyme borreliosis generally occurs in stages.

**Early Lyme Borreliosis:** Although stages may overlap or occur alone, illness may begin with a characteristic skin lesion called erythema migrans (EM) in 60% of cases. This rash appears as a red macule or papule that expands in an annular manner, sometimes with multiple similar lesions. Fever, malaise, fatigue, headache, stiff neck, myalgia, migratory arthralgias, and lymphadenopathy may accompany or precede EM.

**Neurologic Manifestations:** Weeks to months after the onset of early Lyme disease, neurologic abnormalities may develop in untreated patients. The typical pattern is fluctuating meningoencephalitis with superimposed cranial (particularly facial) nerve palsy and peripheral radiculo-neuropathy.

**Cardiac Manifestations:** Within several weeks after onset, about 8% of untreated patients develop cardiac involvement (most commonly fluctuating degrees of atrioventricular block that resolves spontaneously).

**Arthritis:** Weeks to years after the original illness, about 50% of untreated patients develop arthritis. Early involvement typically is manifested by migratory pain, often without swelling. Frank arthritis may develop subsequently with marked swelling and pain in one or more joints, primarily large joints, e.g., the knee.

b. **Differential Diagnosis:**

**Early disease:** Aseptic meningitis, hepatitis, mononucleosis, ehrlichiosis.

**Late disease:** Rheumatic fever, disseminated gonococcal infection, multiple sclerosis, Guillain-Barré syndrome,

Reiter's syndrome, rheumatoid arthritis, oligoarticular form of juvenile rheumatoid arthritis.

c. **Diagnosis:** Based on clinical findings. Serological testing (EIA or IFA) may be useful but lacks sensitivity, especially in early disease. A two-step testing procedure using flagellar protein-based EIA followed by IgM and IgG Western blot of all positive and equivocal specimens is recommended. Culture from biopsy at the outer margins of EM lesion is 90% sensitive. PCR is available from research laboratories.

3. **Incubation:** 7-10 days average, range 3-32 days.

4. **Reservoir:** Wild animals; e.g., *Neotoma* spp. (wood rat) and deer are important in California.

5. **Source:** Infected *Ixodes* species ticks; other arthropods have been found containing *B. burgdorferi*, but their ability to transmit is questionable.

6. **Transmission:** Bite of *Ixodes* tick. 36-48 hours of attachment is usually required for transmission.

7. **Communicability:** Not transmitted from person to person.

8. **Specific Treatment:** Amoxicillin is a good treatment for adults or children with early disease. Doxycycline in adults and phenoxymethyl penicillin for children with early disease resolves illness and reduces the likelihood of later complications. Intravenous penicillin or ceftriaxone is effective for meningitis, late stage, and refractory illness.

## REPORTING PROCEDURES

1. Report any cases or suspected cases within 7 calendar days to ACDC or Morbidity Unit. *California Code of Regulations*, Title 17, Section 2500.

2. **Report Form:** [LYME DISEASE CASE REPORT \(CDPH 8470\)](#).



**3. Epidemiologic Data:**

- a. Travel 30 days prior to onset of erythema migrans or early disease.
- b. History of tick bite.
- c. History of possible exposure to ticks, e.g., hiking in chaparral, dogs with ticks, etc.
- d. Occupational exposure.

**Amount:** 8-10 ml.

**Storage:** Refrigerate.

**Remarks:** Serologic tests for Lyme borreliosis lack sensitivity and are not standardized, so interpretation of test results is difficult. Only confirmed cases are reported to the CDC. Confirmed cases are patients that present with EM with a known exposure, or with clinical findings in addition to appropriate laboratory evidence including: 1) positive culture for *B.burgdorferi* from a clinical specimen, 2) two-tier testing using an EIA or IFA screening test followed by Western blot interpreted using established criteria, or 3) IgG Western blot using established criteria.

The case history form must accompany the specimen(s).

**CONTROL OF CASE, CONTACTS & CARRIERS**

Investigation not required by district staff. Advise ACDC regarding suspect cases; ACDC will supply diagnosing physician with appropriate form or investigate. Initiate investigation within 7 days of notification.

**CASE:** Isolation: None.

**CONTACTS:** No restrictions.

**CARRIERS:** Not applicable.

**PREVENTION-EDUCATION**

1. Use tick repellents.
2. Wear protective clothing in wooded areas.
3. Control ticks on domestic animals.
4. Avoid tick-infested areas when feasible.
5. Check periodically for and carefully remove attached ticks after return from tick-infested areas.

**DIAGNOSTIC PROCEDURES**

**Serology:** Done by commercial laboratories. No longer run at State Laboratory or LAC Public Health Laboratory. Can be run at CDC with prior approval.

**Container:** Serum separator tube

**Laboratory Form:** N/A

**Examination Requested:** IFA, EIA, Western blot.

**Material:** Clotted whole blood.